



CASE REPORT / ПРИКАЗ БОЛЕСНИКА

Histological analysis of bone three months after the treatment of oroantral communication with autologous platelet-rich fibrin – a case series

Kristina Rebić¹, Bojan Janjić¹, Vesna Danilović¹, Vladan Đorđević², Branislav Ilić¹

¹University of Belgrade, School of Dental Medicine, Belgrade, Serbia;

²University of Travnik, Faculty of Pharmacy and Health, Travnik, Federation of Bosnia and Herzegovina, Bosnia and Herzegovina

SUMMARY

Introduction Oroantral communication (OAC) closure may be accompanied by hard and soft tissue defects. Platelet-rich fibrin (PRF) is the second-generation platelet concentrate that can be an effective therapeutic option for the closure of defects up to 5 mm in diameter. The aim of this investigation was to evaluate whether PRF could be a viable therapeutic option for OAC closure and provide sufficient bone quality/quantity for the forthcoming implant placement.

Outlines of cases The case series included eight patients treated with PRF due to the presence of OAC less than 3 mm in diameter. Three months after the surgery, at the time of implant placement, bone samples were harvested and taken for histological analysis. The results demonstrated success in all eight cases, obtaining both hard and soft tissue healing. Histological analysis showed that newly formed bone was present on all histological samples, without visible signs of inflammation and necrosis.

Conclusion PRF could be a viable therapeutic option for OAC closure in specific clinical cases, but future randomized, controlled, clinical studies are required for more conclusive results.

Keywords: bone healing; autologous platelet-rich fibrin; oroantral communication

INTRODUCTION

Surgical closure of the oroantral communication (OAC) may be accompanied by hard and soft tissue defects [1]. Although application of local soft-tissue flaps is still the most utilized technique for OAC closure, recent studies suggest that platelet-rich fibrin (PRF) can be an effective therapeutic option for the closure of defects up to 5 mm in diameter. PRF is associated with minimal postoperative morbidity and allows preservation of adjacent teeth soft tissue structures [2–5]. Additionally, the combination of PRF and bone grafting materials promotes hard tissue healing, obtaining better conditions for future implant placement [3, 4, 5].

PRF is the second-generation platelet concentrate consisting of a three-dimensional polymerized fibrin matrix in a molecular structure. It incorporates blood contents such as leukocytes, erythrocytes, platelets, growth factors, and circulating stem cells [6]. PRF membrane induces tissue regeneration due to the stimulating effects on osteoblast cells, gingival fibroblasts, pulp cells, and periodontal ligament cells [1].

This case series aimed to evaluate whether PRF could be a viable therapeutic option for the OAC closure and provide sufficient bone quality/quantity for the forthcoming implant placement.

REPORT OF CASES

This case series included eight patients treated at the Clinic for Oral Surgery, School of Dental Medicine in Belgrade (six males and two females; aged 21–43 years, mean age 34.6 ± 11.3 years). They were referred to the Clinic due to the presence of OAC and enrolled in the study according to the following inclusion criteria:

- patients in good general health without a history of systemic disease or medication that could interfere with the treatment (ASA1 and ASA2);
- fresh OAC (not more than 24 hours from the tooth extraction);
- without the clinical/radiological signs of maxillary sinusitis;
- long and narrow alveolus of the extracted tooth;
- OAC less than 3 mm in diameter;
- length from the cortical margin of the extracted tooth to the OAC being at least 6 mm;
- clinically compliant patients consent to be enrolled in the study.

The OAC was closed by autologous PRF plugs and membranes, following Choukroun's PRF centrifuge protocol (PRF DUO™, Nice, France) [7]. After the curettage and saline rinsing, wound edges were freshened, and the PRF plug was placed inside the alveolus. PRF membrane was shaped over the site in one layer, and the closure was obtained by interrupted sutures

Received • Примљено:
July 4, 2022

Revised • Ревизија:
April 10, 2023

Accepted • Прихваћено:
April 11, 2023

Online first: May 3, 2023

Correspondence to:

Vladan ĐORĐEVIĆ
University of Travnik
Faculty of Pharmacy and Health
Polje Slavka Gavrančića 17c
72270 Travnik
Federation of Bosnia and
Herzegovina
Bosnia and Herzegovina
vladan.djordjevic@fzf.edu.ba




Figure 1. Platelet-rich fibrin material placed in the alveolar socket



Figure 2. Bone sample inside the trephine burr Ø 3.0 mm

– Figure 1. The follow-up was scheduled, and sutures were removed on the 10th day.

Epithelization and soft tissue healing were uneventful in all eight cases. Three months after the OAC closure, there was a sufficient amount of keratinized gingiva and the cone-beam computed tomography evaluation revealed new bone formation in the area of the previously extracted tooth. Sub-antral height of 6–9 mm (average 7.3 mm) was obtained in all eight cases. The site was reopened, bone samples for histological analysis were harvested (trephine burr; Ø 3.0 mm – Figure 2), and implants were placed. We were using bone level, tapered implants, following the maxilla protocol (avoiding the last sequence drill), and managed to obtain solid implant stability (from 20 Ncm to above).

Bone samples were stored in a 10%-formalin solution for 12–24 hours and then decalcified in a microwave oven (eight cycles of 10 seconds; at 410–430°C for 20 minutes). The material was dehydrated with 70%, 95%, and 99% ethyl alcohol, respectively, and clarified with xylene. Gathered bone fragments were embedded in paraffin blocks, cut into slices (3–4 µm), and stained with Goldner trichrome method. Analysis of samples was performed under Leica Microsystem® optical microscope (Leica Microsystems™ GmbH, Wetzlar, Germany).

The newly formed bone was present in all histological samples, without signs of inflammation and necrosis (Figure 3). Bone trabeculae were surrounded by the loose connective tissue in which no inflammatory infiltrate cells were seen, or their number was minimal. Additionally,




Figure 3. Newly formed bone tissue (Goldner trichrome method, 40 ×)




Figure 4. Newly formed bone tissue with elements of mature and immature bone (Goldner trichrome method, 100 ×)




Figure 5. Osteocytes in the lacunae and osteoid and densely packed osteoblasts (Goldner trichrome method, 400 ×)

intensive bone remodeling was noticed, i.e., the presence of mature (lamellar) bone and immature (fibrous) bone (Figure 4). Haversian canals with concentrically placed bone lamellae (characteristics of osteons) were seen as a sign of mature bone, and collagen-fiber networks as a sign of immature bone. The abundant presence of osteocytes lying in lacunae confirmed bone vitality, and the presence of osteoids covered with a dense layer of osteoblasts was a sign of active osteogenesis (Figure 5).

This report was approved by the institutional ethics committee, and written consent was obtained from the patients for the publication of the report and any accompanying images.

DISCUSSION

Our findings indicate that PRF could be a viable solution for the OAC closure in specific clinical cases. It provides proper hard and soft tissue healing, obtaining sufficient bone quality/quantity. Initially, PRF acts as a boosting agent for soft tissue healing, supporting epithelization. Additionally, it promotes bone formation in the area of the extracted tooth, creating the vital bone, and shortens the healing period. The presence of mature and immature bone is a significant histological sign of intensive bone remodeling.

PRF plug acts as a core for bone healing and the PRF membrane acts as a biological membrane that promotes epithelization. Due to its properties, PRF has proven as the material of choice not only for this indication but in many other clinical studies as well. Ondur et al. [8] showed that the use of PRF for hard and soft tissue healing may have advantages due to its autogenous origin, being cheaper than the collagen membrane. The authors pointed to PRF's ability to release growth factors (TGF- β 1, PDGF- β , VEGF), particularly in the first seven days, and later, up to 28 days. Similarly, Liu et al. [9] promote PRF as a bone grafting material for oral and maxillofacial bone regeneration

procedures as it improves proliferation, migration, differentiation, and mineralization of the cells during bone formation. There is an indication that PRF could also diminish crestal bone resorption after tooth extraction [10], as demonstrated in periimplantitis therapy use [11]. Moreover, the application of PRF, either alone or in combination with another biomaterial, might be effective in reducing time for new bone formation and future implant placement [12, 13].

Although we were successful in all eight cases obtaining hard and soft tissue healing, it would be presumptive to state that PRF could be a universal tool for OAC closure. In this study, we had a strict case selection, requiring fresh OAC with a diameter of up to 3 mm. Post-extraction socket had to be long and narrow, and the distance between the crest and the OAC should be at least 6 mm. However, there is a question if the same healing would be obtained without the use of PRF since we did not have a control group. Additionally, the sample size was small. We did not experience any complications and are considering if the procedure could be applied to larger/shallower defects, along with the use of bone substitute materials.

This case series indicates that PRF could be a viable therapeutic option for the OAC closure providing optimal hard and soft tissue structures for the future implant placement. However, future randomized, controlled studies on larger sample sizes, with control groups, should contribute to more conclusive remarks.

Conflict of interest: None declared.

REFERENCES

- Baek JH, Kim BO, Lee WP. Implant Placement after Closure of Oroantral Communication by Sinus Bone Graft Using a Collagen Barrier Membrane in the Shape of a Pouch: A Case Report and Review of the Literature. *Medicina (Kaunas)*. 2021;57(6):626. [DOI: 10.3390/medicina57060626] [PMID: 34208644]
- Parvini P, Obreja K, Begic A, Schwarz F, Becker J, Sader R, et al. Decision-making in closure of oroantral communication and fistula. *Int J Implant Dent*. 2019;5(1):13. [DOI: 10.1186/s40729-019-0165-7] [PMID: 30931487]
- Salgado-Peralvo AO, Mateos-Moreno MV, Uribarri A, Kewalramani N, Peña-Cardelles JF, Velasco-Ortega E. Treatment of oroantral communication with Platelet-Rich Fibrin: A systematic review. *J Stomatol Oral Maxillofac Surg*. 2022;123(5):e367–e375. [DOI: 10.1016/j.joms.2022.03.014] [PMID: 35318134]
- Khalifaoui M, Eya M, Selmi W, Lamia O, Nabih D. Closure of oroantral communication with platelet-rich fibrin: A case report. *Saudi J Oral Dent Res*. 2021;6(12):566–8. [DOI: 10.36348/sjodr.2021.v06i12.006]
- Mainassara Chekaraou S, Benjelloun L, El Harti K. Management of oro-antral fistula: Two case reports and review. *Ann Med Surg (Lond)*. 2021;69:102817. [DOI: 10.1016/j.jamsu.2021.102817] [PMID: 34527240]
- Niedzielska I, Ciapinski D, Bak M, Niedzielski D. The Assessment of the Usefulness of Platelet-Rich Fibrin in the Healing Process Bone Resorption. *Coatings*. 2022;12(2):247 [DOI: 10.3390/coatings12020247]
- Dohan DM, Choukroun J, Diss A, Dohan SL, Dohan AJ, Mouhyi J, et al. Platelet-rich fibrin (PRF): a second-generation platelet concentrate. Part I: technological concepts and evolution. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2006;101(3):e37–44. [DOI: 10.1016/j.tripleo.2005.07.008] [PMID: 16504849]
- Ondur E, Bolukbasi Balcioglu N, Soluk Tekkesin M, Guzel O, Ersanli S. Effects of Platelet-Rich Fibrin on Hard Tissue Healing: A Histomorphometric Crossover Trial in Sheep. *Materials (Basel)*. 2020;13(7):1695. [DOI: 10.3390/ma13071695] [PMID: 32260464]
- Liu Y, Sun X, Yu J, Wang J, Zhai P, Chen S, et al. Platelet-Rich Fibrin as a Bone Graft Material in Oral and Maxillofacial Bone Regeneration: Classification and Summary for Better Application. *Biomed Res Int*. 2019;2019:3295756. [DOI: 10.1155/2019/3295756] [PMID: 31886202]
- Al-Badran A, Bierbaum S, Wolf-Brandstetter C. Does the Choice of Preparation Protocol for Platelet-Rich Fibrin Have Consequences for Healing and Alveolar Ridge Preservation After Tooth Extraction? A Meta-Analysis. *J Oral Maxillofac Surg*. 2023;81(5):602–21. [DOI: 10.1016/j.joms.2023.01.004] [PMID: 36736375]
- Sun G, Cao L, Li H. Effects of platelet-rich fibrin combined with guided bone regeneration in the reconstruction of peri-implantitis bone defect. *Am J Transl Res*. 2021;13(7):8397–402. [PMID: 34377334]
- Otero AIP, Fernandes JCH, Borges T, Nassani L, Castilho RM, Fernandes GVO. Sinus Lift Associated with Leucocyte-Platelet-Rich Fibrin (Second Generation) for Bone Gain: A Systematic Review. *J Clin Med*. 2022;11(7):1888. [DOI: 10.3390/jcm11071888] [PMID: 35407494]
- Guan S, Xiao T, Bai J, Ning C, Zhang X, Yang L, et al. Clinical application of platelet-rich fibrin to enhance dental implant stability: A systematic review and meta-analysis. *Heliyon*. 2023;9(2):e13196. [DOI: 10.1016/j.heliyon.2023.e13196] [PMID: 36785817]

Хистолошка анализа кости три месеца после реконструисања ороантралне комуникације аутологним фибрином богатим тромбоцитима – серија болесника

Кристина Ребић¹, Бојан Јањић¹, Весна Даниловић¹, Владан Ђорђевић², Бранислав Илић¹

¹Универзитет у Београду, Стоматолошки факултет, Београд, Србија;

²Универзитет у Травнику, Фармацеутско-здравствени факултет, Травник, Федерација Босне и Херцеговине, Босна и Херцеговина

САЖЕТАК

Увод Хируршко затварање ороантралних комуникација (ОАК) може бити праћено стварањем коштаних или мекоткивних дефеката. Аутологни фибрин богат тромбоцитима (ФБТ) јесте тромбоцитни концентрат друге генерације ефикасан у реконструкцији ОАК дијаметра до 5 mm.

Циљ истраживања је био да испита ефикасност ФБТ у реконструкцији ОАК и утврди да ли ће његова примена обезбедити адекватну коштану подлогу за будућу уградњу имплантата.

Приказ болесника Истраживање је обухватило осам пацијената код којих је ОАК била мања од 3 mm и реконструисана

применом ФБТ. Три месеца после хируршког захвата, приликом уградње имплантата, са места ОАК узети су узорци кости ради хистолошке анализе. Мекоткивно нарастање је било успешно код свих испитаника. Резултати свих узорака показали су присуство новоформиране здраве кости, без знакова запаљења и некрозе.

Закључак ФБТ се може користити за реконструкцију ОАК у специфичним клиничким индикацијама. Ипак, неопходно је спровести рандомизоване, контролисане клиничке студије пре доношења јасних препорука.

Кључне речи: коштано нарастање; аутологни фибрин богат тромбоцитима; ороантрална комуникација